

**TITLE: TRANSCUTANEOUS CARDIAC
PACING COMPETENCY**

EMS Policy No. 2547

SKILLS: TRANSCUTANEOUS CARDIAC PACING (TCP)

Name: _____ **Date:** _____ **Cert #** _____

Transcutaneous pacing is used for short intervals as a bridge until transvenous pacing can be initiated or until the underlying cause of the bradyarrhythmia (e.g., hyperkalemia, drug overdose) can be reversed.

A. Assessment/Treatment Indicators:

1. Indicated for adult patients with hemodynamically unstable bradycardia.
2. Is authorized as a standing order for paramedics in treating adult patients with unstable bradycardia. Hemodynamically unstable bradycardia means a patient with a BP < 90, related to a bradycardic rhythm (HR <60) with serious signs and symptoms related to heart rate, (i.e.: chest pain, SOB, ALOC, shock, pulmonary congestion, CHF).
3. TCP should not be delayed for hemodynamically unstable bradycardia patients while waiting for IV access or for atropine to take effect.

B. Contraindications: TCP is not authorized for use on patients less than 15 years of age. Not authorized for hypothermic patients because the bradycardia is usually a physiologic response to the body temperature.

<u>Performance Criteria</u>		<u>Met Initials</u>	<u>Not Met Initials</u>	<u>Comments</u>
1.	Explain procedure (to patient).			
2.	State indications and contraindications for pacing.			
3.	Apply pre-gelled adhesive pacing pads to chest wall according to manufacturer's recommendations.			
4.	Apply ECG electrodes			
5.	Confirm rhythm.			
6.	Activate pacing device per manufacturer's instructions.			
7.	Set heart rate			
8.	Increases output until capture occurs (and increase output 10% above threshold)			
9.	Confirms capture by correlating QRS spike with pulses. Reassesses BP and LOC. Increases rate prn (not to exceed 100) if patient remains hypotensive and symptomatic from inadequate perfusion.			

Effective: January 1, 2009

Page 1 of 2

Revised:

Supersedes:

Approved: Signature on File
Medical Director

Signature on File
EMS Administrator

**TITLE: TRANSCUTANEOUS CARDIAC
PACING COMPETENCY**

EMS Policy No. **2547**

<u>Performance Criteria</u>		<u>Met Initials</u>	<u>Not Met Initials</u>	<u>Comments</u>
10.	Determines what the lowest threshold response and maintains output control at this level. NOTE: Any movement of patient may increase the capture threshold response; subsequently, the output may have to be adjusted to compensate for this.			
11.	Provides patient sedation/pain relief prn.			
12.	Continue monitoring. Contact base hospital for further orders if patient symptoms are not resolving (consideration for dopamine, further alteration of pacer settings or if further sedation/pain control orders required).			

VALIDATOR'S SIGNATURE	ALS PROVIDER'S SIGNATURE	DATE

Effective: January 1, 2009

Page 2 of 2

Revised:

Supersedes:

Approved: Signature on File
Medical Director

Signature on File
EMS Administrator