San Joaquin County Emergency Medical Services Agency



BLS Abdominal Emergencies

AUTHORITY: Health and Safety Code, Division 2.5, Sections 1797.60, 1797.80, 1797.197, 1797.197a, 1797.204, 1797.220, 1798; California Code of Regulations, Title 22, Chapter 2, Sections 100062, 100063, 100064

INFORMATION NEEDED:

Discomfort: location, quality, severity, onset of symptoms, duration, aggravation, alleviation. Associated symptoms – nausea, emesis, diarrhea, fever, diaphoresis, vertigo, "heart burn". Gastro-intestinal – time and description of last meal, time of last bowel movement, signs of

blood in stool.

Gynecological: date of last menstrual period, possible pregnancy, history of vaginal bleeding. Medical history: surgery, related diagnosis (infection, hepatitis, stones, etc.), medication

(OTC and prescribed), self administered remedies (baking soda, Epsom salts, enemas).

OBJECTIVE FINDINGS:

General appearance – level of distress, skin color, diaphoresis. Abdominal tenderness – guarding, rigidity, distention, rebound. Pulsating masses (aneurysm). Quality of femoral pulses.

TREATMENT:

- 1. Primary Survey ensure ABC's.
- 2. Place patient in position of comfort or supine with legs elevated if patient is hypotensive.
- 3. Monitor SpO2.
- 4. Administer oxygen for shortness of breath or signs of hypoxia: Oxygen 10-15 L/min via non-rebreathing mask. Patients with ineffective respirations: support ventilations with BVM and airway.
- 5. Give patient nothing by mouth.
- 6. Confirm ambulance transport is responding.
- 7. Secondary Survey and Routine Medical Care.
- Note: Completion of a thorough secondary exam and patient history are essential to identify potential cardiac involvement or early signs of shock.

Effective:April 1, 2020Supersedes:July 1, 2007

Approved: <u>Signature on file</u> Medical Director

Page 1 of 1

Signature on file EMS Administrator