San Joaquin County Emergency Medical Services Agency



BLS Head - Neck- Facial Trauma

AUTHORITY: Health and Safety Code, Division 2.5, Sections 1797.60, 1797.80, 1797.197, 1797.197a, 1797.204, 1797.220, 1798; California Code of Regulations, Title 22, Chapter 2, Sections 100062, 100063, 100064

INFORMATION NEEDED:

Mechanism of injury.

Medical history: cardiovascular problems, diabetes, or seizure disorder.

OBJECTIVE FINDINGS:

Check for DCAP-BTLS (Deformity, Contusion/Crepitus, Abrasion, Puncture, Bleeding, Tenderness, Laceration, Swelling).

Signs or airway obstruction: stridor, abnormal voice, difficulty breathing.

Glasgow Coma Score.

Neurological impairment or focal deficit (paralysis, weakness).

Eyes/vision: pupil equality and reactivity, eye tracking, impaired vision (double vision, stars).

TREATMENT:

- 1. Primary Survey ABC's.
- 2. Monitor SpO2.
- 3. Administer oxygen for shortness of breath or signs of hypoxia: Oxygen 10-15 L/min via non-rebreathing mask. Patients with ineffective respirations: support ventilations with BVM and appropriate airway adjunct.
- 4. Provide spinal motion restriction if indicated by mechanism of injury and patient assessment.
- 5. Control external bleeding with direct pressure and hemostatic dressings as necessary.
- 6. Stabilize impaled objects with bulky dressings.
- 7. Apply cold packs to reduce pain and decrease soft tissue swelling.
- 8. Following specific treatment: Secondary Survey and Routine Medical Care.

Specific treatment:

- 9. Eye injury Apply dressing as appropriate, loosely cover affected and unaffected eye.
- 10. Tooth injury keep avulsed teeth in saline soaked gauze (or commercial tooth saver kit) and transport with patient.
- 11. Mandible fracture splint with cravat or bandage.

Effective: April 1, 2020 Page 1 of 2

Supersedes: January 1, 2012

Approved: Signature on file Signature on file

Medical Director EMS Administrator

Notes:

- 1) All patients with a period of unconsciousness should be transported to an emergency department for evaluation.
- 2) Continually monitor Glasgow Coma Score and observe for fluid drainage from ear or nose.

Effective: April 1, 2020 Page 2 of 2

Approved: Signature on file Signature on file

Supersedes: January 1, 2012

Medical Director EMS Administrator